

Desirability

The essentials of imprecise-probability theory
from a different perspective

Erik Quaeghebeur

SYSTeMS & FUM Research Groups, Ghent University

Outline

Sets of desirable gambles

- Basic concepts

- Coherent sets of desirable gambles

- Assessments, avoiding partial loss & extensions

- Marginally desirable gambles & friends

- Updating sets of desirable gambles

Partial preference orders

Lower and upper previsions

- Moving from sets of desirable gambles to previsions

- Assessments, avoiding sure loss, natural extension & coherence

- Linear previsions & credal sets

- Updating lower previsions

Basic concepts: context & assumptions

Possibility space Ω e.g., outcomes experiment

Subject uncertain about outcome experiment

Goal model the subject's beliefs & use this model for reasoning

Gambles payoff depends on outcome,
bounded real-valued function on Ω ,
set of gambles \mathcal{L}_Ω


Utility linear and precise

Interpretation behavioral

Basic concepts: (desirable) gambles

$$\Omega := \{a, b\}$$

$$f \in \mathcal{L}_\Omega$$



Gamble f desirable when the subject accepts the transaction

- (i) the experiment's outcome ω is determined
- (ii) the subject's capital is changed by $f(\omega)$


Subject's belief model set of desirable gambles

Rationality criteria for sets of desirable gambles: passive

To be *coherent*, a set of desirable gambles \mathcal{R} must satisfy

Accepting partial gains: $\mathcal{L}_\Omega^+ \subseteq \mathcal{R}$,

Avoiding partial loss: $\mathcal{L}_\Omega^- \cap \mathcal{R} = \emptyset$.



The zero gamble is on the fence.

Rationality criteria for sets of desirable gambles: active

To be *coherent*, a set of desirable gambles \mathcal{R} must also satisfy


Positive scaling: $\lambda > 0, f \in \mathcal{R} \Rightarrow \lambda f \in \mathcal{R}$,

Addition: $f, g \in \mathcal{R} \Rightarrow f + g \in \mathcal{R}$.

A coherent set of desirable gambles \mathcal{R} is a *convex cone* including \mathcal{L}_Ω^+ and excluding \mathcal{L}_Ω^- :

$$\lambda \mathcal{R} = \mathcal{R}$$

$$\mathcal{R} + \mathcal{R} = \mathcal{R}$$



Some people who have worked with cones

C.A.B. Smith 1961 implicitly, using an open cone of 'exchange vectors'

P.M. Williams 1974 all main ideas present, uses the word 'acceptable'
builds on B. de Finetti's work



P. Walley 1991 focus on conceptual ease of updating,
builds on the above two

S. Moral 2000 epistemic irrelevance,
later work also with collaborators

G. de Cooman & E. Miranda 2007 symmetry

Assessments


An **assessment** is a set \mathcal{A} of gambles judged desirable by the subject.



Conical hull operator coni all finite strictly positive linear combinations of elements of its argument set

Avoiding partial loss

An assessment \mathcal{A} incurs partial loss iff $\text{coni}(\mathcal{A}) \cap \mathcal{L}_\Omega^- \neq \emptyset$.
No coherent set of desirable gambles will encompass it then.




An assessment \mathcal{A} avoids partial loss (and thus sure loss) iff

$$\text{coni}(\mathcal{A}) \cap \mathcal{L}_\Omega^- = \emptyset$$

Least and maximally committal extensions

An assessment that avoids partial loss can have a continuum of encompassing coherent sets of desirable gambles \mathcal{R}




Maximal ones are not always (closed or open) half-spaces!

Natural extension

The least committal extension of an assessment \mathcal{A} is called its *natural extension*


$$\mathcal{E}(\mathcal{A}) = \text{coni}(\mathcal{A} \cup \mathcal{L}_\Omega^+).$$



The natural extension $\mathcal{E}(\mathcal{A})$ is coherent iff \mathcal{A} avoids partial loss.

Natural extension as an intersection

The set of cones is closed under intersections.





The natural extension of an assessment that avoids partial loss is the intersection of all

- ▶ coherent sets of desirable gambles that encompass it
- ▶ maximal elements of the coherent sets of desirable gambles that encompass it
- ▶ a minimal set of maximal elements of the coherent sets of desirable gambles that encompass it

Marginally desirable gambles

The set of marginally desirable gambles of \mathcal{R} is



$$\begin{aligned}\mathcal{G}_{\mathcal{R}} &:= \{f - \sup\{\alpha \in \mathbb{R} \mid f - \alpha \in \mathcal{R}\} \mid f \in \mathcal{R}\} \\ &= \text{bd}(\mathcal{R}) \quad \text{when } \mathcal{R} \text{ coherent.}\end{aligned}$$



Marginally desirable gambles & friends

The set of marginally desirable gambles of \mathcal{R} is

$$\begin{aligned}\mathcal{G}_{\mathcal{R}} &:= \{f - \sup\{\alpha \in \mathbb{R} \mid f - \alpha \in \mathcal{R}\} \mid f \in \mathcal{R}\} \\ &= \text{bd}(\mathcal{R}) \quad \text{when } \mathcal{R} \text{ coherent.}\end{aligned}$$



The set of almost desirable gambles is $\text{cl}(\mathcal{R})$, the closure of \mathcal{R}

The set of strictly desirable gambles is $\text{int}(\mathcal{R}) \cup \mathcal{L}_{\Omega}^{+}$,
the interior of \mathcal{R} and gambles from \mathcal{L}_{Ω}^{+}



Updating a set of desirable gambles

Indicator I_C of an event $C \subseteq \Omega$ is the gamble $I_C(\omega) := \begin{cases} 1, & \omega \in C, \\ 0, & \text{otherwise.} \end{cases}$

Updated set of desirable gambles of a set of desirable gambles \mathcal{R} on an event C :







$$\mathcal{R}_C := \{f_C \mid f \in \mathcal{R}; f = fI_C\} \subseteq \mathcal{L}_C$$

$$\Omega \supset C = \{a, b\}$$



Updating a coherent set of desirable gambles results in a coherent updated set of desirable gambles.

Updated sets of desirable gambles & marginally desirable gambles



Things left unmentioned

- ▶ Sets of desirable gambles relative to a linear subspace of \mathcal{L}_Ω
- ▶ Transformations of Ω and sets of desirable gambles
- ▶ The issue of the zero gamble
- ▶ Combining different sets of desirable gambles

Outline

Sets of desirable gambles

- Basic concepts

- Coherent sets of desirable gambles

- Assessments, avoiding partial loss & extensions

- Marginally desirable gambles & friends

- Updating sets of desirable gambles

Partial preference orders

Lower and upper previsions



- Moving from sets of desirable gambles to previsions

- Assessments, avoiding sure loss, natural extension & coherence

- Linear previsions & credal sets

- Updating lower previsions


Difference of gambles




Comparing & ordering gambles

Partial preference order \succeq associated to
a coherent set of desirable gambles \mathcal{R} :

$$f \succeq g \Leftrightarrow f - g \in \mathcal{R}$$





$f \not\succeq g$ and $g \not\succeq f$



$f \succeq g$ and $g \not\succeq f$


Difference of gambles




Comparing & ordering gambles

Partial preference order \succeq associated to
a coherent set of desirable gambles \mathcal{R} :

$$f \succeq g \Leftrightarrow f - g \in \mathcal{R}$$



$f \stackrel{?}{\succeq} g$ and $g \not\stackrel{?}{\succeq} f$



$f \stackrel{?}{\succeq} g$ and $g \stackrel{?}{\succeq} f$

Outline

Sets of desirable gambles

- Basic concepts

- Coherent sets of desirable gambles

- Assessments, avoiding partial loss & extensions

- Marginally desirable gambles & friends

- Updating sets of desirable gambles

Partial preference orders

Lower and upper previsions


- Moving from sets of desirable gambles to previsions

- Assessments, avoiding sure loss, natural extension & coherence

- Linear previsions & credal sets

- Updating lower previsions


Constant gambles



- ▶ The set of constant gambles can be linearly ordered (trivially so).
- ▶ What happens if we compare a gamble f with all constant gambles?
- ▶ Use constant gambles for buying and selling gambles.

From (marginally) desirable gambles to previsions


Lower and upper previsions \underline{P} and \bar{P} corresponding to a coherent set of desirable gambles \mathcal{R} :



$$\begin{aligned}\underline{P}(f) &:= \sup\{\alpha \in \mathbb{R} \mid f \geq \alpha\} = \sup\{\alpha \in \mathbb{R} \mid f - \alpha \in \mathcal{R}\} \\ &= \alpha \text{ such that } f - \alpha \in \mathcal{G}_{\mathcal{R}}\end{aligned}$$

From (marginally) desirable gambles to previsions

Lower and upper previsions \underline{P} and \bar{P} corresponding to a coherent set of desirable gambles \mathcal{R} :




$$\begin{aligned}\bar{P}(f) &:= \inf\{\beta \in \mathbb{R} \mid \beta \geq f\} = \inf\{\beta \in \mathbb{R} \mid \beta - f \in \mathcal{R}\} \\ &= -\underline{P}(-f)\end{aligned}$$

Lower and upper probabilities

Lower and upper probabilities of the event A :

$$\underline{P}(A) := \underline{P}(I_A) \quad \bar{P}(A) := \bar{P}(I_A) = 1 - \underline{P}(\Omega \setminus A)$$




Lower and upper probabilities are less expressive than lower and upper previsions, which are less expressive than sets of desirable gambles.

From previsions to assessments

What is the assessment corresponding to a lower prevision \underline{P} on $\mathcal{K} \subseteq \mathcal{L}_\Omega$?

The set of marginally desirable gambles of \underline{P} is

$$\mathcal{G}_{\underline{P}} := \{G_{\underline{P}}(f) \mid f \in \mathcal{K}\}, \quad \text{with } G_{\underline{P}}(f) = f - \underline{P}(f).$$




$$\mathcal{A}_{\underline{P}} := \mathcal{G}_{\underline{P}} + \mathbb{R}_{>0} = \{g + \alpha \mid g \in \mathcal{G}_{\underline{P}}; \alpha > 0\}$$

Avoiding sure loss

A lower prevision \underline{P} on $\mathcal{K} \subseteq \mathcal{L}_\Omega$ avoids sure loss iff

- ▶ $\mathcal{A}_{\underline{P}}$ avoids sure loss, or, as it is an open set, iff $\mathcal{A}_{\underline{P}}$ avoids partial loss

$$\text{coni}(\mathcal{A}_{\underline{P}}) \cap \mathcal{L}_\Omega^- = \emptyset \quad \Leftrightarrow \quad \forall g \in \text{coni}(\mathcal{G}_{\underline{P}}) : \sup\{g\} \geq 0$$





$$\mathcal{A}_{\underline{P}} := \mathcal{G}_{\underline{P}} + \mathbb{R}_{>0} = \{g + \alpha \mid g \in \mathcal{G}_{\underline{P}}; \alpha > 0\}$$

Natural extension of a lower prevision

The natural extension \underline{E} of a lower prevision \underline{P} on $\mathcal{K} \subseteq \mathcal{L}_\Omega$ is

$$\begin{aligned}\underline{E}(f) &:= \sup\{\alpha \in \mathbb{R} \mid f - \alpha \in \mathcal{E}(\underline{P})\}, \quad \text{with } \mathcal{E}(\underline{P}) := \mathcal{E}(\mathcal{A}_{\underline{P}}), \\ &= \sup\{\alpha \in \mathbb{R} \mid f - \alpha \geq g; g \in \text{coni}(\mathcal{G}_{\underline{P}})\}.\end{aligned}$$





Coherence of a lower prevision

A lower prevision \underline{P} on $\mathcal{K} \subseteq \mathcal{L}_\Omega$ is *coherent* iff

$$\forall f \in \mathcal{K} : \underline{P}(f) = \underline{E}(f)$$

\Leftrightarrow


$$\forall f \in \mathcal{G}_{\underline{P}} : \forall g \in \text{coni}(\mathcal{G}_{\underline{P}}) : \sup\{g - f\} \geq 0.$$



Linear previsions & hyperplanes


A linear prevision P corresponds to a set of marginally desirable gambles \mathcal{G}_P that is a hyperplane:

$$P(G_P(f)) = P(f - P(f)) = 0 = P(P(f) - f) = P(G_P(-f)).$$



Maximally committal extensions & credal sets

The natural extension $\mathcal{E}(\underline{P})$ of a lower prevision \underline{P} that avoids sure loss is the intersection of the set of maximally committal extensions.



The credal set $\mathcal{M}_{\underline{P}}$ is the closed convex set of linear previsions

$$\mathcal{M}_{\underline{P}} := \{P \text{ on } \mathcal{L}_{\Omega} \mid P \geq \underline{P}\}$$



The natural extension \underline{E} of \underline{P} can then be written as

$$\underline{E}(f) = \min\{P \in \mathcal{M}_{\underline{P}} \mid P(f)\} = \min\{P \in \text{ext } \mathcal{M}_{\underline{P}} \mid P(f)\}.$$

Updating a coherent lower prevision

Updated lower prevision of a set of desirable gambles \mathcal{R} or a coherent lower prevision \underline{P} on \mathcal{L}_Ω (then $\mathcal{R} := \mathcal{E}(\underline{P})$):

$$\begin{aligned}\underline{P}(g|C) &:= \sup\{\alpha \in \mathbb{R} \mid g - \alpha \in \mathcal{R}_C\} \\ &= \sup\{\alpha \in \mathbb{R} \mid (g - \alpha)I_C \in \mathcal{R}\}\end{aligned}$$



The updated lower prevision $\underline{P}(\cdot|C)$ is a coherent lower prevision on \mathcal{L}_C .

Updating with natural and regular extension

Natural extension or the generalized Bayes's rule; alternative formulations:

$$\begin{aligned}\underline{P}(g|C) &= \mu \in \mathbb{R} \text{ for which } \underline{P}((g - \mu)I_C) = 0 \\ &= \min\{P(gI_C)/P(C) \mid P \in \mathcal{M}_{\underline{P}}\}\end{aligned}$$

whenever $\underline{P}(C) > 0$, otherwise $\underline{P}(\cdot|C)$ is vacuous.

Updating with natural and regular extension

Natural extension or the generalized Bayes's rule; alternative formulations:

$$\begin{aligned}\underline{P}(g|C) &= \mu \in \mathbb{R} \text{ for which } \underline{P}((g - \mu)I_C) = 0 \\ &= \min\{P(gI_C)/P(C) \mid P \in \mathcal{M}_{\underline{P}}\}\end{aligned}$$

whenever $\underline{P}(C) > 0$, otherwise $\underline{P}(\cdot|C)$ is vacuous.

Regular extension

$$\begin{aligned}\underline{R}(g|C) &= \max\{\mu \in \mathbb{R} \mid \underline{P}((g - \mu)I_C) \geq 0\} \\ &= \inf\{P(gI_C)/P(C) \mid P \in \mathcal{M}_{\underline{P}}; P(C) > 0\}\end{aligned}$$

whenever $\bar{P}(C) > 0$, otherwise $\underline{R}(\cdot|C)$ is vacuous.

Updating with natural and regular extension

Natural extension or the generalized Bayes's rule; alternative formulations:




$$\begin{aligned}\underline{P}(g|C) &= \mu \in \mathbb{R} \text{ for which } \underline{P}((g - \mu)I_C) = 0 \\ &= \min\{P(gI_C)/P(C) \mid P \in \mathcal{M}_{\underline{P}}\}\end{aligned}$$

whenever $\underline{P}(C) > 0$, otherwise $\underline{P}(\cdot|C)$ is vacuous.

Regular extension

$$\begin{aligned}\underline{R}(g|C) &= \max\{\mu \in \mathbb{R} \mid \underline{P}((g - \mu)I_C) \geq 0\} \\ &= \inf\{P(gI_C)/P(C) \mid P \in \mathcal{M}_{\underline{P}}; P(C) > 0\}\end{aligned}$$

whenever $\bar{P}(C) > 0$, otherwise $\underline{R}(\cdot|C)$ is vacuous.



Updating on events of lower probability zero with natural and regular extension




Consider a coherent set of desirable gambles \mathcal{R}

- ▶ $\underline{Q}(\cdot|C)$ is the lower prevision associated to \mathcal{R}_C
- ▶ \underline{P} is the lower prevision associated to \mathcal{R}
and $\bar{P}(C) > \underline{P}(C) = 0$, so $\underline{P}(\cdot|C)$ is vacuous
- ▶ $\underline{R}(\cdot|C)$ is the regular extension of \underline{P}

Updating on events of lower probability zero with natural and regular extension

Consider a coherent set of desirable gambles \mathcal{R}

- ▶ $\underline{Q}(\cdot|C)$ is the lower prevision associated to \mathcal{R}_C
- ▶ \underline{P} is the lower prevision associated to \mathcal{R} and $\bar{P}(C) > \underline{P}(C) = 0$, so $\underline{P}(\cdot|C)$ is vacuous
- ▶ $\underline{R}(\cdot|C)$ is the regular extension of \underline{P}




$$\inf\{g\} = \underline{P}(g|C) \leq \underline{Q}(g|C) \leq \underline{R}(g|C)$$

Updating on events of upper probability zero with natural and regular extension

Consider a coherent set of desirable gambles \mathcal{R}

- ▶ $\underline{Q}(\cdot|C)$ is the lower prevision associated to \mathcal{R}_C
- ▶ \underline{P} is the lower prevision associated to \mathcal{R}
and $\bar{P}(C) = \underline{P}(C) = 0$, so $\underline{P}(\cdot|C)$ is vacuous
- ▶ $\underline{R}(\cdot|C)$ is the regular extension of \underline{P} ; it is also vacuous



$$\inf\{g\} = \underline{P}(g|C) = \underline{R}(g|C) \leq \underline{Q}(g|C)$$

Outline

Sets of desirable gambles

- Basic concepts

- Coherent sets of desirable gambles

- Assessments, avoiding partial loss & extensions

- Marginally desirable gambles & friends

- Updating sets of desirable gambles

Partial preference orders

Lower and upper previsions

- Moving from sets of desirable gambles to previsions

- Assessments, avoiding sure loss, natural extension & coherence

- Linear previsions & credal sets

- Updating lower previsions

Advocating desirability

- ▶ A conceptually simple basis for imprecise-probability theory
- ▶ Intuitive and pedagogical due to its geometric nature
- ▶ Updating is elegant
- ▶ Holds promise for studying structural assessments

Combining updated sets of desirable gambles

- ▶ Given an assessment \mathcal{A}_A for each event A in a set of events \mathcal{A} of Ω
- ▶ These can be combined in an assessment on Ω :

$$\mathcal{A} := \bigcup_{A \in \mathcal{A}} \{gI_A \mid g \in \mathcal{A}_A\}$$

Combining updated sets of desirable gambles

- ▶ Given an assessent \mathcal{A}_A for each event A in a set of events \mathcal{A} of Ω
- ▶ These can be combined in an assessment on Ω :

$$\mathcal{A} := \bigcup_{A \in \mathcal{A}} \{g I_A \mid g \in \mathcal{A}_A\}$$

- ▶ Given an assessent \mathcal{A}_B for each event B in a partition \mathcal{B} of Ω
- ▶ These can be combined in an assessment on Ω :

$$\mathcal{A} := \left\{ \sum_{B \in \mathcal{B}} g^B I_B \mid g^B \in \mathcal{A}_B; B \in \mathcal{B} \right\}$$

Combining updated sets of desirable gambles

- ▶ Given an assessent \mathcal{A}_A for each event A in a set of events \mathcal{A} of Ω
- ▶ These can be combined in an assessment on Ω :

$$\mathcal{A} := \bigcup_{A \in \mathcal{A}} \{g I_A \mid g \in \mathcal{A}_A\}$$

- ▶ Given an assessent \mathcal{A}_B for each event B in a partition \mathcal{B} of Ω
- ▶ These can be combined in an assessment on Ω :

$$\mathcal{A} := \left\{ \sum_{B \in \mathcal{B}} g^B I_B \mid g^B \in \mathcal{A}_B; B \in \mathcal{B} \right\}$$

- ▶ What about infinite partitions?
- ▶ \mathcal{B} -Conglomerability:

$$\forall B \in \mathcal{B} : f_B \in \mathcal{A}_B \Rightarrow f \in \mathcal{A}$$

(‘Full conglomerability’ if it must hold for all partitions \mathcal{B} of Ω .)

Conditional previsions & marginal gambles

Conditional lower prevision $\underline{P}(\cdot|\mathcal{B})$ for a partition \mathcal{B} of Ω :

$$\underline{P}(f|\mathcal{B})(\omega) := \underline{P}(f|B) = \underline{P}(f_B|B) \text{ when } \omega \in B$$


It is a *piecewise constant gamble*, constant on $B \in \mathcal{B}$.

Conditional previsions & marginal gambles

Conditional lower prevision $\underline{P}(\cdot|\mathcal{B})$ for a partition \mathcal{B} of Ω :

$$\underline{P}(f|\mathcal{B})(\omega) := \underline{P}(f|B) = \underline{P}(f_B|B) \text{ when } \omega \in B$$

It is a *piecewise constant gamble*, constant on $B \in \mathcal{B}$.



$$\Omega := A \cup B$$

$$\mathcal{B} := \{A, B\}$$

$$C := \{a, b\}$$

$$a \in A$$

$$b \in B$$

References I



G. de Cooman and E. Miranda.

Symmetry of models versus models of symmetry.

In *Probability and Inference: Essays in Honor of Henry E. Kyburg, Jr.*, pages 67–149. King's College Publications, 2007.



B. de Finetti.

Teoria delle Probabilità.

Einaudi, Turin, 1970.



B. de Finetti.

Theory of Probability: A Critical Introductory Treatment.

John Wiley & Sons, Chichester, 1974–1975.

English translation of [de Finetti(1970)], two volumes.

References II



S. Moral.

Epistemic irrelevance on sets of desirable gambles.

In ISIPTA '01 – Proceedings of the Second International Symposium on Imprecise Probabilities and Their Applications, pages 247–254.

Shaker Publishing, Maastricht, 2000.



S. Moral.

Epistemic irrelevance on sets of desirable gambles.

Annals of Mathematics and Artificial Intelligence, 45:197–214, 2005.



E. Quaeghebeur.

Learning from samples using coherent lower previsions.

PhD thesis, Ghent University, 2009.



C. A. B. Smith.

Consistency in statistical inference and decision.

Journal of the Royal Statistical Society, Series A, 23:1–37, 1961.

References III



P. Walley.

Statistical Reasoning with Imprecise Probabilities.

Chapman and Hall, London, 1991.



P. Walley.

Towards a unified theory of imprecise probability.

International Journal of Approximate Reasoning, 24:125–148, 2000.



P. M. Williams.

Indeterminate probabilities.

In *Formal Methods in the Methodology of Empirical Sciences*, pages 229–246. 1974.



P. M. Williams.

Notes on conditional previsions.

Technical report, School of Mathematical and Physical Science, University of Sussex, UK, 1975.

Revised journal version: [Williams 2007].

References IV



P. M. Williams.

Coherence, strict coherence and zero probabilities.

In Proceedings of the Fifth International Congress on Logic, Methodology and Philosophy of Science, volume VI, pages 29–33. 1975.



P. M. Williams.

Notes on conditional previsions.

International Journal of Approximate Reasoning, 44:366–383, 2007.
Revised journal version of [Williams 1975a].